

Whole-Genome Draft Sequences of Six Commensal Fecal and Six Mastitis-Associated *Escherichia coli* Strains of Bovine Origin

Andreas Leimbach,^{a,b,c}  Anja Poehlein,^b Anika Witten,^d Olga Wellnitz,^e Nahum Shpigel,^f Wolfram Petzl,^g Holm Zerbe,^g Rolf Daniel,^b  Ulrich Dobrindt^{a,c}

Institute for Hygiene, University of Münster, Münster, Germany^a; Department of Genomic and Applied Microbiology, Göttingen Genomics Laboratory, Institute of Microbiology and Genetics, Georg-August-University of Göttingen, Göttingen, Germany^b; Institute for Molecular Infection Biology, Julius-Maximilians-University of Würzburg, Würzburg, Germany^c; Institute for Human Genetics, University of Münster, Münster, Germany^d; Veterinary Physiology, Vetsuisse Faculty University of Bern, Bern, Switzerland^e; Koret School of Veterinary Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel^f; Clinic for Ruminants, Ludwig-Maximilians-University of Munich, Oberschleißheim, Germany^g

The bovine gastrointestinal tract is a natural reservoir for commensal and pathogenic *Escherichia coli* strains with the ability to cause mastitis. Here, we report the whole-genome sequences of six *E. coli* isolates from acute mastitis cases and six *E. coli* isolates from the feces of udder-healthy cows.

Received 6 June 2016 Accepted 10 June 2016 Published 28 July 2016

Citation Leimbach A, Poehlein A, Witten A, Wellnitz O, Shpigel N, Petzl W, Zerbe H, Daniel R, Dobrindt U. 2016. Whole-genome draft sequences of six commensal fecal and six mastitis-associated *Escherichia coli* strains of bovine origin. *Genome Announc* 4(4):e00753-16. doi:10.1128/genomeA.00753-16.

Copyright © 2016 Leimbach et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ulrich Dobrindt, dobrindt@uni-muenster.de.

Although bovine intramammary infections with *Escherichia coli* mostly lead to an acute onset of mastitis, they can also result in a persistent infection of the udder with alternating sub-clinical or clinical periods (1). Additionally, no common virulence factor subset of mastitis-causing *E. coli* strains has been identified in previous studies (2).

To investigate the genomic potential of *E. coli* isolated from bovine mastitis, several draft genomes (3–5), as well as two complete genomes (6), have been published thus far. However, only two recent genomic *E. coli* mastitis studies included one commensal bovine isolate (7, 8). Because cows are a natural reservoir not only for pathogenic but also for commensal *E. coli* of high phylogenetic and genotypic diversity (2), we present here the draft genomes of six *E. coli* strains isolated from serous udder exudate of mastitis-afflicted cows and six *E. coli* strains isolated from the feces of udder-healthy cows (Table 1).

All genomes were sequenced with an Illumina HiScan SQ sequencer with Nextera XT chemistry (Illumina, San Diego,

CA, USA) for library preparation and a 101-bp paired-end sequencing run. Raw reads were quality controlled with FastQC version 0.11.2 (<http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc>). Low-quality reads and adapter contaminations were trimmed with Cutadapt version 1.6 (9). All reads were randomly subsampled to an approximate 70-fold coverage for each strain with seqtk version 1.0-r32 (<https://github.com/lh3/seqtk>). Subsequently, the reads were *de novo* assembled with SPAdes version 3.1.1 (10). Assembly statistics were evaluated with QUAST version 2.3 (11), resulting in 59 to 290 contigs >500 bp and genome sizes ranging from 4,765,494 to 5,459,392 bp (Table 1).

The strains were classified evenly into phylogroups A or B1, regardless of isolation source, through the assignment of sequence types (ST) with *e. coli*_mlst version 0.3 (https://github.com/aleimba/bac-genomics-scripts/tree/master/ecoli_mlst) (12). The most prominent sequence type is ST10, but most of the strains were not closely phylogenetically related.

All genomes were annotated with Prokka version 1.9 (13) with

TABLE 1 Genome features and assembly metrics of the 12 *E. coli* whole-genome sequences

Strain	ECOR phylogroup (ST)	Source of isolation	Genome size (bp)	No. of contigs >500 bp	<i>N</i> ₅₀ (bp)	No. of CDSs ^a	Accession no.
131/07	A (ST10)	Udder acute mastitis	5,459,392	270	79,414	5,123	JXUH000000000
2772a	B1 (ST156)	Udder acute mastitis	4,949,901	93	163,837	4,621	LCVG000000000
3234/A	A (ST10)	Udder acute mastitis	5,482,981	290	95,923	5,211	LCVH000000000
MPEC4839	A (ST10)	Udder acute mastitis	4,866,885	124	133,521	4,502	JYHP000000000
MPEC4969	B1 (ST1125)	Udder acute mastitis	4,833,611	130	103,834	4,468	JYHQ000000000
RiKo 2299/09	B1 (ST448)	Healthy cow feces	4,954,750	129	114,991	4,587	JYKB000000000
RiKo 2305/09	B1 (ST410)	Healthy cow feces	4,806,931	123	129,952	4,429	JYPB000000000
RiKo 2308/09	A (ST167)	Healthy cow feces	5,112,873	186	83,735	4,685	LCVI000000000
RiKo 2331/09	B1 (ST1614)	Healthy cow feces	4,765,494	59	224,192	4,350	LCVJ000000000
RiKo 2340/09	A (ST167)	Healthy cow feces	5,024,854	204	82,522	4,568	LAGW000000000
RiKo 2351/09	B1 (ST88)	Healthy cow feces	5,297,190	252	102,610	4,931	LAUC000000000
UVM2	A (ST1091)	Udder acute mastitis	4,926,170	149	86,033	4,614	LAUD000000000

^a CDS, coding sequence.

E. coli 1303 (CP009166 to CP009169) or *E. coli* ECC-1470 (CP010344 to CP010345) as references for annotation for either the ECOR phylogroup A or B1 genomes, respectively. tRNAs were predicted with tRNAscan-SE version 1.3.1 (14). Additionally, the annotations were manually curated with Proteinortho version 5.11 (15), po2anno version 0.2 (<https://github.com/aleimba/bac-genomics-scripts/tree/master/po2anno>), ACT version 13.0.0 (16), and *E. coli* strains 1303 and ECC-1470 as references. Finally, tbl2tab version 0.2 (<https://github.com/aleimba/bac-genomics-scripts/tree/master/tbl2tab>) and Artemis version 16.0.0 (17) were used to refine the annotations after querying the Virulence Factors Database (18) and the ResFinder version 2.1 (19), VirulenceFinder version 1.2 (20), and SerotypeFinder version 1.0 (21) databases. In summary, between 4,350 and 5,211 coding DNA sequences were identified in the genomes with 3 to 7 rRNAs and 68 to 83 tRNAs.

The genome sequences in this study will serve as a useful resource for future comparative studies of *E. coli* strains associated with bovine mastitis in relationship to commensal strains and for the identification of potential virulence factors.

Nucleotide sequence accession numbers. These whole-genome shotgun projects have been deposited at DDBJ/EMBL/GenBank under the accession numbers listed in Table 1. The versions described here are the first versions.

ACKNOWLEDGMENT

We thank David E. Kerr (University of Vermont, Burlington) for providing *E. coli* mastitis isolate UVM2.

FUNDING INFORMATION

This work, including the efforts of Ulrich Dobrindt, was funded by Deutsche Forschungsgemeinschaft (DFG) (DO 789/3-1 and DO 789/4-1). This work, including the efforts of Andreas Leimbach, was funded by Deutsche Forschungsgemeinschaft (DFG) (DO 789/3-1). This work, including the efforts of Wolfram Petz and Holm Zerbe, was funded by Deutsche Forschungsgemeinschaft (DFG) (FOR 585).

REFERENCES

- Zadoks RN, Middleton JR, McDougall S, Katholm J, Schukken YH. 2011. Molecular epidemiology of mastitis pathogens of dairy cattle and comparative relevance to humans. *J Mammary Gland Biol Neoplasia* 16: 357–372. <http://dx.doi.org/10.1007/s10911-011-9236-y>.
- Blum SE, Leitner G. 2013. Genotyping and virulence factors assessment of bovine mastitis *Escherichia coli*. *Vet Microbiol* 163:305–312. <http://dx.doi.org/10.1016/j.vetmic.2012.12.037>.
- Blum S, Sela N, Heller ED, Sela S, Leitner G. 2012. Genome analysis of bovine-mastitis-associated *Escherichia coli* O32:H37 strain P4. *J Bacteriol* 194:3732. <http://dx.doi.org/10.1128/JB.00535-12>.
- Kempf F, Loux V, Germon P. 2015. Genome sequences of two bovine mastitis-causing *Escherichia coli* strains. *Genome Announc* 3(2):e00259-15. <http://dx.doi.org/10.1128/genomeA.00259-15>.
- Richards VP, Lefebvre T, Pavinski Bitar PD, Dogan B, Simpson KW, Schukken YH, Stanhope MJ. 2015. Genome based phylogeny and comparative genomic analysis of intra-mammary pathogenic *Escherichia coli*. *PLoS One* 10:e0119799. <http://dx.doi.org/10.1371/journal.pone.0119799>.
- Leimbach A, Poehlein A, Witten A, Scheutz F, Schukken Y, Daniel R, Dobrindt U. 2015. Complete genome sequences of *Escherichia coli* strains 1303 and ECC-1470 isolated from bovine mastitis. *Genome Announc* 3(2):e00182-15. <http://dx.doi.org/10.1128/genomeA.00182-15>.
- Blum SE, Heller ED, Sela S, Elad D, Edery N, Leitner G. 2015. Genomic and phenomic study of mammary pathogenic *Escherichia coli*. *PLoS One* 10:e0136387. <http://dx.doi.org/10.1371/journal.pone.0136387>.
- Kempf F, Slugocki C, Blum SE, Leitner G, Germon P. 2016. Genomic comparative study of bovine mastitis *Escherichia coli*. *PLoS One* 11: e0147954. <http://dx.doi.org/10.1371/journal.pone.0147954>.
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10. <http://dx.doi.org/10.14806/ej.17.1.200>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <http://dx.doi.org/10.1093/bioinformatics/btt086>.
- Wirth T, Falush D, Lan R, Colles F, Mensa P, Wieler LH, Karch H, Reeves PR, Maiden MC, Ochman H, Achtman M. 2006. Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* 60: 1136–1151. <http://dx.doi.org/10.1111/j.1365-2958.2006.05172.x>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25: 955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
- Lechner M, Findeiss S, Steiner L, Marz M, Stadler PF, Prohaska SJ. 2011. Proteinortho: detection of (co-)orthologs in large-scale analysis. *BMC Bioinformatics* 12:124. <http://dx.doi.org/10.1186/1471-2105-12-124>.
- Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis comparison tool. *Bioinformatics* 21: 3422–3423. <http://dx.doi.org/10.1093/bioinformatics/bti553>.
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945. <http://dx.doi.org/10.1093/bioinformatics/16.10.944>.
- Chen L, Xiong Z, Sun L, Yang J, Jin Q. 2012. VFDB 2012 update: toward the genetic diversity and molecular evolution of bacterial virulence factors. *Nucleic Acids Res* 40:D641–D645. <http://dx.doi.org/10.1093/nar/gkr989>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.
- Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <http://dx.doi.org/10.1128/JCM.03617-13>.
- Joensen KG, Tetzschner AM, Iguchi A, Aarestrup FM, Scheutz F. 2015. Rapid and easy *in silico* serotyping of *Escherichia coli* isolates by use of whole-genome sequencing data. *J Clin Microbiol* 53:2410–2426. <http://dx.doi.org/10.1128/JCM.00008-15>.